RESULTS GUARANTEED†

Kills Over 250 Types of Weeds! as listed

Spectracide®

WEED STOP® FOR LAWNS

PLUS CRABGRASS KILLER

CONCENTRATE

Herbicida concentrado para céspedes y contra mano de cangrejo

Visible Results in 8 Hours

Treats up to 5,000 sq ft

Won't Harm Lawns When Used as Directed

MIX WITH WATER™

17-12655

ACTIVE INGREDIENTS:	
2,4-D, dimethylamine salt	3.74%
Quinclorac	1.79%
Dicamba, dimethylamine salt	0.43%
Sulfentrazone	0.22%
OTHER INGREDIENTS	93.82%
TOTAL	100.00%

THIS PRODUCT CONTAINS: 0.264 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 3.11%, 0.152 lb 3,7-dichloro-8-quinolinecarboxylic acid per gallon or 1.79%, 0.030 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 0.36%, 0.019 lb N-[2,4-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]phenyl]methanesulfonamide per gallon or 0.22%. Isomer Specific by AOAC Method.

KEEP OUT OF REACH OF CHILDREN

CAUTION See back booklet for additional precautionary statements.

†For terms of guarantee, see back panel.

MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS

PRECAUCIÓN Consulte el folleto en la parte posterior para conocer las precauciones adicionales.

Net 32 fl oz (1 qt/ 946 mL)

- **Para COMENZAR a rociar:** Conecte la manguera al rociador. Extienda la manguera de jardín hasta el sector de césped a tratar que se encuentre más alejado. Gire la punta de la boquilla del rociador para dirigir el rocío hacia arriba o abajo. Abra el agua y apriete el gatillo o gire la perilla para comenzar.
- **DESPUÉS de rociar:** El producto o concentrado sin usar se puede volver a verter en el envase original. Mantenga el ajuste del selector y limpie el rociador enjuagando la jarra del rociador con agua durante un minuto.

Para mayores detalles sobre el uso apropiado de su rociador, revise las indicaciones que recibió al adquirirlo.

PARA ROCIADORES PARA CONCENTRADO MÁS AGUA O ROCIADORES CON APLICADOR TIPO MANGUERA CON PREMEZCLA

Para rociadores que requieren el agregado de agua y concentrado antes de la aplicación:

1. A partir de la Tabla 3, determine la cantidad de concentrado necesaria para tratar todo el césped. Vierta el concentrado dentro de la jarra del rociador hasta el nivel necesario de "onza líquida."
2. Agregue la cantidad adecuada de agua hasta el nivel de "galones." Vuelva a poner la tapa del rociador en la jarra y agite bien antes de rociar.
3. Conecte el rociador a la manguera. Extienda la manguera hasta la zona más alejada que se vaya a tratar y retroceda hacia el grifo. Aplique uniformemente al césped.

Tabla 3. Dosis de aplicación para césped con rociadores con aplicador tipo manguera que requieren premezcla.

Área que se va a tratar, pies cuadrados	Pasto azul de Kentucky, festuca spp., ryegrass perenne, césped zoysia, pasto Bermuda** y pasto búfalo	Marca/nivel de galón en la jarra del rociador
1,250 pies cuadrados	8 onzas líquidas	3 galones
1,875 pies cuadrados	12 onzas líquidas	5 galones
2,500 pies cuadrados	16 onzas líquidas	6 galones
3,750 pies cuadrados	24 onzas líquidas	9 galones

**La aplicación al pasto Bermuda puede hacer que se torne amarillo o se decolore temporalmente, pero puede esperar su total recuperación. No aplique al pasto Bermuda cuando las temperaturas diurnas superen los 85°F.

Ejemplo para pasto azul de Kentucky: Para tratar 3,750 pies cuadrados de césped, vierta 24 onzas líquidas de producto en la jarra del rociador. Agregue agua hasta la marca de 9 galones y aplique uniformemente sobre 3,750 pies cuadrados de césped.

Para mayores detalles sobre el uso apropiado de su rociador, revise las indicaciones que recibió al adquirirlo.

CONSEJOS SOBRE EL USO

Las condiciones ambientales templadas y de humedad adecuada mejoran el rendimiento de este producto. Las aplicaciones se deben llevar a cabo en malezas que se encuentran en la etapa de desarrollo activo. Aplique cuando las temperaturas diurnas se encuentren entre 40°F y 90°F. Evite aplicar este producto cuando los céspedes estén sometidos a estrés ya que puede ocasionar daño.

Para áreas recién sembradas:

- No aplique en el plazo de 4 semanas después de la germinación de las plantas de semillero de pasto azul de Kentucky, mezclas de festuca de hoja fina y ryegrass perenne. Aplique este producto a céspedes recién cultivados cuando estén bien establecidos (aproximadamente 6 semanas después de la germinación de las plantas de semillero) o después del tercer corte.

Para áreas recién cubiertas de césped, de propagación vegetativa ("sprigging") o con "plugs":

- La aplicación de este producto a pastos recién cubiertos, de propagación vegetativa ("sprigging') o con "plugs" se debe demorar hasta 3 ó 4 semanas después de operaciones de cobertura de césped, propagación vegetativa o colocación de "plugs."

Siembra:

- Las especies de césped que se indican en esta etiqueta se pueden sembrar en las áreas tratadas 4 semanas después de la aplicación de este producto.

Riego:

- No aplique este producto inmediatamente antes de la lluvia o el riego. Para obtener mejores resultados, no irrigue ni riegue el césped por 24 horas después de la aplicación.
- Si las condiciones son secas, se recomienda programar la irrigación o el riego 24 horas antes y 24 horas después de la aplicación.

Corte:

- Retrase el corte 2 días antes y hasta 2 días después de la aplicación de este producto. Más estrés ocasionado por la altura baja del corte puede aumentar la posibilidad del daño al césped.
- Los recortes de los primeros 3 cortes se deben dejar en el área tratada. No use los recortes provenientes del área tratada como mantillo ni abono orgánicos alrededor de flores, plantas ornamentales, árboles ni en huertas.

PROHIBICIONES:

- No trate cuando la temperatura del aire sea superior a 85°F porque se puede dañar el césped.
- La aplicación de dosis superiores a las que se especifican en esta etiqueta puede causar daño al césped.
- No aplique este producto a cultivos de alimentos (jardines y hortalizas), cultivos de forraje ni plantas ornamentales, que incluyen flores, árboles, arbustos, setos, plantas ornamentales leñosas, ni en cubresuelos establecidos en plantaciones ornamentales y otras plantas que no sean el objetivo.
- No rocíe las raíces expuestas de árboles y plantas ornamentales.

ALMACENAMIENTO Y ELIMINACIÓN

Almacenamiento del pesticida: Guárdelo en el envase original. No contamine el agua, ni los alimentos para personas ni animales por medio del almacenamiento o la eliminación.

Eliminación:

Si está vacío: No vuelva a usar este envase. Tírelo en la basura, u ofrézcalo para reciclar, si existe esa opción. **Si está parcialmente vacío:** Llame a su servicio de eliminación de desechos más cercano para obtener instrucciones de eliminación. Nunca tire el producto restante por ningún drenaje interno ni externo.

PRECAUCIONES

Riesgos para los seres humanos y los animales domésticos

PRECAUCIÓN. Causa irritación moderada a los ojos. Evite el contacto con los ojos o la ropa. Nocivo si se ingiere. Cuando utilice este producto, use camisa de manga larga, pantalones largos, calcetines, zapatos y guantes de goma/hule.

Después de manipular el producto, lávese bien con agua y jabón. Después de utilizar este producto, enjuague los guantes antes de quitárselos, quítese la ropa y lávela por separado antes de volver a usarla y rápidamente lávese bien las manos y la piel que estuvo expuesta, con agua y jabón. Quítese la ropa mojada y báñese en regadera tan pronto como sea posible.

Primeros Auxilios	
Si entra en contacto con los ojos:	• Mantenga los ojos abiertos y enjuague con agua lentamente, con cuidado, durante 15-20 minutos. Si hay lentes de contacto, retírelos después de los primeros 5 minutos, luego continúe enjuagando los ojos. • Llame a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento.
Si se ingiere:	• Llame inmediatamente a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento. • Si la persona puede tragar, haga que beba poco a poco un vaso de agua. • No induzca el vómito a menos que así lo indique el centro de control de envenenamientos o el médico. • No administre nada a una persona que haya perdido el conocimiento.
Cuando llame a un centro de control de envenenamientos o a un médico, o intente obtener tratamiento, tenga a la mano el envase o la etiqueta del producto. También puede llamar al 1-800-917-5438 para obtener información acerca del tratamiento médico de urgencia.	

RIESGOS PARA EL MEDIO AMBIENTE

Este pesticida es tóxico para los peces y los invertebrados acuáticos. No lo aplique directamente al agua, a zonas donde haya aguas superficiales ó áreas intermareales por debajo de la marca de promedio correspondiente al nivel más alto del agua excepto cuando lo indiquen las etiquetas apropiadas. El arrastre y el escurrimiento pueden resultar peligrosos para los organismos acuáticos de las zonas cercanas al sitio de aplicación. No contamine el agua cuando elimine las aguas de lavado del equipo ni los residuos del enjuague.

Este producto químico tiene propiedades y características relacionadas con sustancias químicas detectadas en aguas subterráneas. El uso de este producto químico en zonas de suelos permeables, particularmente donde la capa freática es poco profunda, puede ocasionar la contaminación del agua subterránea. La aplicación alrededor de un tanque de agua o un pozo puede contaminar el agua potable o subterránea.

Spectracide

STOP READ ENTIRE LABEL BEFORE USE. ALTO. LEA TODA LA ETIQUETA ANTES DE USAR.

PEEL HERE RESEALABLE LABEL
ABRA AQUÍ ETIQUETA RESELLABLE

WEED STOP® FOR LAWNS

PLUS CRABGRASS KILLER

CONCENTRATE

Herbicida concentrado para céspedes y contra mano de cangrejo

Kills on Contact

Treats up to 5,000 sq ft

Rainproof in as Little as 3 Hours

Kills Weeds, Including Crabgrass – Not the Lawn

Kills Weed Roots!

Kills 250+ Types of Weeds as Listed

PRODUCT FACTS

WHAT IT DOES:
Kills major broadleaf and troublesome grass weeds including dandelion, chickweed and clover as well as crabgrass, foxtail and yellow nutsedge. (See inside for complete list).

WHERE TO USE:
Lawns: Kills the weeds, not the lawn.

WHEN TO USE:
Apply when daytime temperatures are between 45°F and 90°F. **Do not apply to Bermudagrass when daytime temperatures exceed 85°F.**

QUESTIONS & COMMENTS:
Call 1-800-917-5438 or visit our website at **www.spectracide.com**

†GUARANTEED RESULTS OR YOUR MONEY BACK

If you are not satisfied with this product for any reason, send your written request for a refund to the address below, together with the original dated store receipt showing the price you paid. Please allow 6-8 weeks for processing.

Application to Bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

NOTA: El comprador asume toda la responsabilidad en caso de no seguir las instrucciones sobre seguridad y modo de empleo.

Distributed by Spectrum Group
Div. of United Industries Corp.
PO Box 142642, St. Louis, MO 63114-0642
EPA Reg. No. 9688-268-8845
EPA Est. Nos. 39754-WI-1 (R), 9688-MO-1 (U), 79533-WI-1 (BT), 39609-MO-1 (BM)
Circled letter is first letter of lot number.
17-12654 © 2008 UIC

PROOF OF PURCHASE

FPO UPC
0 71121 95702 3

ALTO LEA TODA LA ETIQUETA ANTES DE USAR.

MODO DE EMPLEO

Se considera una violación a la ley federal usar este producto con otros fines que los indicados en la etiqueta. *Sólo para uso residencial al aire libre.*

AGITE BIEN ANTES DE USAR.

- No aplique este producto de modo que entre en contacto con personas o mascotas, ya sea directamente o por arrastre. Mantenga a las personas y las mascotas fuera del área durante la aplicación.
- No permita que las personas ni las mascotas entren en el área tratada hasta que los rocíos se hayan secado.

INFORMACIÓN GENERAL:

El herbicida concentrado para céspedes y contra mano de cangrejo Spectracide® Weed Stop® es un herbicida selectivo que controla malezas de hoja ancha y malezas herbosas que germinaron en céspedes establecidos. Algunos de los síntomas previstos de las malezas de hoja ancha son enrojecimiento de las hojas, doblamiento y torsión de hojas y tallo, clorosis y necrosis. Los síntomas previstos en las malezas herbosas consisten en atrofia, clorosis, enrojecimiento gradual y necrosis.

DÓNDE USAR

Este producto está destinado al uso por parte de los propietarios en céspedes residenciales. Este producto sólo debe usar en las siguientes especies de césped.

Césped de temporada fría	**Césped de temporada cálida**
Pasto azul de Kentucky	Pasto Bermuda**
Ryegrass perenne	Césped zoysia
Festuca spp., incluso festuca alta, roja y de hoja fina	Pasto búfalo

**La aplicación al pasto Bermuda puede provocar que el césped se vuelva amarillo o se decolore temporalmente pero se puede esperar una recuperación completa. No lo aplique al pasto Bermuda cuando las temperaturas diurnas superan los 85°F.

Este producto **NO** se debe usar en céspedes que contengan agróstide, césped San Agustín, césped bahía, césped ciempiés o césped alfombra. No lo use en céspedes dichondra ni en céspedes con tréboles o legumbres deseables, en huertas o frutas, ni en jardines con plantas ornamentales (flores, árboles, cubresuelos, setos, lechos de flores o plantas y arbustos).

CUÁNDO USARLO

Para obtener mejores resultados, aplique este producto a malezas que se encuentren en la etapa de desarrollo activo durante las etapas iniciales. Aplique en primavera u otoño cuando las temperaturas diurnas se encuentren entre 45°F y 90°F. Las aplicaciones en el otoño pueden controlar las malezas anuales y bienales que de lo contrario vuelven a germinar en la primavera siguiente. Si hay muchas malezas herbosas haga una segunda aplicación con la misma dosis, 14 días después de la primera aplicación.

MANO DE CANGREJO (grande y suave):

- **Primavera:** Para obtener mejores resultados, aplique este producto desde la emergencia del pasto mano de cangrejo hasta la etapa de seis hojas (dos retoños) o antes de que el pasto mano de cangrejo tenga 3 a 4 pulgadas de altura. Este período generalmente comienza o corresponde a la primera vez que usted observa mano de cangrejo en su césped. Haga dos aplicaciones con la misma dosis con un intervalo de 14 días para poblaciones densas.

CUÁNTO USAR

Vea la tabla siguiente para calcular la cantidad correcta de producto a usar para el área que se va a tratar.
La dosis máxima de aplicación es de 6½ onzas líquidas de producto cada 1,000 pies cuadrados por aplicación (0.6 lb. de equivalente ácido de 2,4-D por acre por aplicación).
La cantidad máxima de aplicaciones al voleo se limita a dos por año.
La dosis estacional máxima es de 13 onzas líquidas de producto cada 1,000 pies cuadrados (1.2 lb. de equivalente ácido de 2,4-D por acre), sin incluir los tratamientos localizados.

CÁLCULOS DE MEDICIÓN: El área total de césped en pies cuadrados = largo x ancho.

MEDIDAS EQUIVALENTES:

1 onza líquida = 2 cdas. soperas = 6 cdtas	4 onzas líquidas = ½ taza	16 onzas líquidas = 1 pinta (pt)
2 onza líquida = 4 cdas. soperas	8 onzas líquidas = 1 taza	32 onzas líquidas = 1 cuarto de galón

I. TRATAMIENTO LOCALIZADO

Para lograr un mejor control, cuando trate áreas pequeñas o malezas individuales, use un rociador a presión. Mezcle la cantidad de producto indicada con 1 galón de agua cada 500 pies cuadrados que se vayan a tratar. Agite bien antes de rociar. Rocíe completamente las malezas que sean el objetivo. Pueden ser necesarios ajustes de la boquilla para administrar de manera uniforme las gotitas pequeñas del rocío en las malezas que son el objetivo.

Cantidad de producto que se debe diluir en 1 galón de agua necesaria para tratar 500 pies cuadrados de césped con rociadores a presión.

Césped	Cantidad de producto — onza líquida (cdas. soperas)	Área tratada aproximada
Pasto azul de Kentucky, festuca spp., ryegrass perenne, césped zoysia, pasto Bermuda** y pasto búfalo	5 onzas líquidas ó 10 cdas. soperas	500 pies cuadrados

**La aplicación al pasto Bermuda puede hacer que se torne amarillo o se decolore temporalmente, pero puede esperar su total recuperación. No aplique al pasto Bermuda cuando las temperaturas diurnas superen los 85°F.

II. TRATAMIENTOS DE TODO EL CÉSPED

Tres pasos fáciles para mezclar y aplicar con rociadores manuales:

1. Calcule (mida) el área total de césped que se va a tratar. En el caso de áreas rectangulares, multiplique la longitud del área por el ancho para obtener la superficie total en pies cuadrados. Prepare suficiente cantidad de solución de rocío para tratar el área calculada.
2. Prepare la mezcla para rociar de acuerdo a la Tabla 1. Mida la cantidad adecuada de producto para tratar 1,000 pies cuadrados. Si es necesario tratar más o menos que 1,000 pies cuadrados, ajuste la cantidad de producto como se indica en la Tabla 1.
3. Agite la solución de rocío antes y durante la aplicación. Use un tipo de rocío que descargue gotas pequeñas pero que no forme niebla.

TRATAMIENTOS AL VOLEO:

ROCIADORES A PRESIÓN

Tabla 1. Cantidad de producto para tratar 1,000 pies cuadrados de césped con rociadores a presión.

Césped	Cantidad de producto a usar	Área a tratar
Pasto azul de Kentucky, festuca spp., ryegrass perenne, césped zoysia, pasto Bermuda** y pasto búfalo	6.4 onzas líquidas	1,000 pies cuadrados
	12.8 onzas líquidas	2,000 pies cuadrados
	32 onzas líquidas	5,000 pies cuadrados

**La aplicación al pasto Bermuda puede hacer que se torne amarillo o se decolore temporalmente, pero puede esperar su total recuperación. No aplique al pasto Bermuda cuando las temperaturas diurnas superen los 85°F.

PARA ROCIADORES CON SELECTOR CON APLICADOR TIPO MANGUERA SIN PREMEZCLA

1. Utilizando la Tabla 2 siguiente, determine la cantidad de producto que se debe aplicar cada 1,000 pies cuadrados de césped.
2. Llene la jarra del rociador con la cantidad de concentrado adecuada para cubrir toda el área que se va a tratar. NO agregue agua a la jarra del rociador.
3. Ajuste el selector del rociador en la posición "Dosis por galón" indicada en la Tabla 2.
4. Conecte el rociador a la manguera de jardín. Extienda la manguera hasta la zona más alejada que se vaya a tratar y retroceda hacia el grifo. Aplique uniformemente al césped.
5. El producto sin usar se puede volver a poner en el envase original al terminar.

Tabla 2. Dosis de aplicación para césped con rociadores con aplicador tipo manguera.

Césped	Dosis por galón o ajustes del selector	Cantidad de herbicida aplicada cada 1,000 pies cuadrados‡
Pasto azul de Kentucky, festuca spp., ryegrass perenne, césped zoysia, pasto Bermuda** y pasto búfalo	5 cdas. soperas ó 15 cdas. ó 2.5 onzas líquidas	6.4 onzas líquidas

‡Se supone que el rociador descarga aproximadamente 2½ galones cada 1,000 pies cuadrados rociados
**La aplicación al pasto Bermuda puede hacer que se torne amarillo o se decolore temporalmente, pero puede esperar su total recuperación. No aplique al pasto Bermuda cuando las temperaturas diurnas superen los 85°F.

Ejemplo para pasto azul de Kentucky: Para tratar 3,000 pies cuadrados de césped, mida 20 onzas líquidas de producto en la jarra del rociador. No agregue agua. **Ajuste el selector en 5 cdas. soperas ó 15 cdtas.** y aplique uniformemente sobre 3,000 pies cuadrados de césped.

Amount of product to be diluted in 1 gal water required to treat 500 sq ft of lawn with pressure sprayers.

Turfgrass	Amount of product fl oz (Tbsp)	Approximate treated area
Kentucky bluegrass, Fescue spp., Perennial ryegrass, Zoysiagrass, Bermudagrass** and Buffalograss	5 fl oz or 10 Tbsp	500 sq ft

**Application to Bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to Bermudagrass when daytime temperatures exceed 85˚F.

II. ENTIRE LAWN TREATMENTS

Three easy steps to mix and apply with hand-operated sprayers:

1. Calculate (measure) the total lawn area to be treated. For rectangular areas, multiply the length of the area by the width to obtain total square feet. Prepare enough spray solution to treat the area calculated.
2. Prepare the spray mixture according to Table 1. Measure the proper amount of product to treat 1,000 sq ft. If more or less than 1,000 sq ft is to be treated, adjust the amount of product as shown in Table 1.
3. Shake the spray solution before and during application. Use a spray pattern that gives small droplets without any fine mist.

BROADCAST TREATMENTS:

PRESSURE SPRAYERS

Table 1. Amount of product to treat 1,000 sq ft of lawn with pressure sprayers.

Turfgrass	Amount of product to use	Area to be treated
Kentucky bluegrass, Fescue spp., Perennial ryegrass, Zoysiagrass, Bermudagrass** and Buffalograss	6.4 fl oz	1,000 sq ft
	12.8 fl oz	2,000 sq ft
	32 fl oz	5,000 sq ft

**Application to Bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to Bermudagrass when daytime temperatures exceed 85˚F.

FOR DIAL-IN HOSE-END SPRAYERS WITH NO PRE-MIXING

1. From Table 2 below, determine amount of this product to be applied for each 1,000 sq ft of lawn.
2. Fill sprayer jar with enough concentrate to cover the total area to be treated. DO NOT add water to the sprayer jar.
3. Set dial on sprayer to the "Rate per Gallon" setting shown in Table 2.
4. Attach sprayer to garden hose. Extend hose to the farthest area of lawn to be treated and work toward faucet. Apply evenly to lawn.
5. Unused product can be returned to the original container when completed.

Table 2. Application rates for turfgrass with hose-end sprayers.

Turfgrass	Rate per gallon or dial settings	Amount of herbicide applied per 1,000 sq ft‡
Kentucky bluegrass, Fescue spp., Perennial ryegrass, Zoysiagrass, Bermudagrass** and Buffalograss	5 Tbsp or 15 tsp or 2.5 fl oz	6.4 fl oz

‡This assumes the sprayer delivers approximately 2½ gal per 1,000 sq ft sprayed

**Application to Bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to Bermudagrass when daytime temperatures exceed 85˚F.

Example for Kentucky bluegrass: To treat 3,000 sq ft of lawn, measure 20 fl oz of product into sprayer jar. Do not add water. **Set dial on 5 Tbsp or 15 tsp** and apply uniformly on 3,000 sq ft of turfgrass.

- **To BEGIN spraying:** Attach hose to sprayer. Extend the garden hose to the farthest section of lawn to be treated. Rotate sprayer nozzle tip to direct spray up or down. Turn on water and squeeze trigger or rotate on/off knob to begin.
- **AFTER spraying:** Unused product or concentrate can be poured back into the original container. Keep the dial setting and clean the sprayer by rinsing the sprayer jar with water for one minute.

Check the directions you received when you purchased your sprayer for further details on the proper use of your dial sprayer.

FOR CONCENTRATE-PLUS-WATER SPRAYERS OR HOSE-END SPRAYERS WITH PRE-MIXING

For sprayers that require the addition of water and concentrate before application:

1. From Table 3, determine the amount of concentrate required to treat the entire lawn. Pour concentrate into sprayer jar to the required "fl oz" level.
2. Add the appropriate amount of water to the "gallons" level. Replace sprayer top on jar and shake well before spraying.
3. Attach sprayer to hose. Extend hose to the farthest area of lawn to be treated and work toward faucet. Apply evenly to lawn.

Table 3. Application rates for turfgrass with hose-end sprayers that require pre-mixing.

Area to be treated, sq ft	Kentucky bluegrass, Fescue spp., Perennial ryegrass, Zoysiagrass, Bermudagrass** and Buffalograss	Gallon mark/level on sprayer jar
1,250 sq ft	8 fl oz	3 gal
1,875 sq ft	12 fl oz	5 gal
2,500 sq ft	16 fl oz	6 gal
3,750 sq ft	24 fl oz	9 gal

**Application to Bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to Bermudagrass when daytime temperatures exceed 85˚F.

Example for Kentucky bluegrass: To treat 3,750 sq ft of lawn, pour 24 fl oz of product into sprayer jar. Add water to the 9 gal mark, and apply uniformly on 3,750 sq ft of turfgrass.

Check the directions you received when you purchased your sprayer for further details on the proper use of your sprayer.

USE TIPS

Adequate moisture and mild environmental conditions enhance the performance of this product. Applications should be made to actively growing weeds. Apply when daytime temperatures are between 40°F and 90°F. Avoid applications of this product when turfgrasses are under stress since injury may result.

For newly seeded areas:

- Do not apply within 4 weeks after seedling emergence of Kentucky bluegrass, fine fescue blends and perennial ryegrass. Apply this product to newly seeded grasses when well-established (approximately 6 weeks after seedling emergence) or after the third mowing.

For newly sodded, sprigged or plugged areas:

- The application of this product to newly sodded, sprigged or plugged grasses should be delayed until 3 to 4 weeks after the sodding, sprigging or plugging operations.

Seeding:

- Turf species listed on this label can be seeded into the treated areas at 4 weeks after the application of this product.

Irrigation:

- Do not apply this product immediately before rainfall or irrigation. For best results, do not irrigate or water the turfgrass within 24 hours after application.
- If dry conditions exist, a scheduled irrigation or watering 24 hours before and 24 hours after application is recommended.

Mowing:

- Delay mowing 2 days before and until 2 days after the application of this product. Additional stress from low mowing heights may increase the possibility of turf injury.
- Clippings from the first 3 mowings should be left on the treated area. Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

PROHIBITIONS:

- Do not treat when air temperatures exceed 85°F as damage to grass may occur.
- Application at rates above specified on this label may cause injury to turfgrass.
- Do not apply this product to food crops (gardens and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants.
- Do not spray exposed roots of ornamentals and trees.

STORAGE AND DISPOSAL

Pesticide Storage: Store in original container. Do not contaminate water, food or feed by storage or disposal.

Disposal:

If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION. Causes moderate eye irritation. Avoid contact with eyes or clothing. Harmful if swallowed. When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves.

Wash thoroughly with soap and water after handling. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

First Aid	
If in eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first five minutes, then continue rinsing eye. • Call a Poison Control Center or doctor for treatment advice.
If swallowed:	• Call a Poison Control Center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a Poison Control Center or doctor. • Do not give anything to an unconscious person.
Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment. You may also contact 1-800-917-5438 for emergency medical treatment advice.	

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high-water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

BROADLEAF WEEDS CONTROLLED/MALEZAS DE HOJA ANCHA CONTROLADAS:

Amaranth, purple
Amaranth, slender
Aster, heath
Aster, purple
Aster, white heath
Aster, white prairie
Artichoke
Austrian fieldcress
Bedstraw, catchweed
Bedstraw, northern
Bedstraw, pineywoods
Bedstraw, smooth
Beggartícks, hairy
Betony, Florida
Bindweed, field
Bindweed, hedge
Bitter wintercress
Bittercress, hairy
Bitterweed
Black-eyed Susan
Black medic
Blood flower milkweed
Blue lettuce
Brassbuttons
Bristly oxtongue
Broomweed
Buckhorn
Bullnettle
Burclover
Burdock
Burweed, lawn (spurweed)
Buttercup, bulbous
Buttercup, creeping
Buttonweed, common (poorjoe)
Buttonweed, Virginia
Carolina geranium (wild)
Carpetweed
Catnip
Catsear, spotted (false dandelion)
Chamber bitter
Chickweed, common
Chickweed, mouseear
Chickweed, sticky
Chicory
Cinquefoil
Clover, alsike
Clover, crimson
Clover, hop
Clover, rabbit foot
Clover, red
Clover, strawberry
Clover, white
Clover, white sweet
Clover, yellow sweet
Cockle
Cocklebur
Cocklebur, oriental
Cocklebur, spiny
Common mullein
Corn spurry
Creeping beggarweed
Creeping Jenny (creeping Charlie)
Cupid shaving brush
Cudweed
Daisy, English
Daisy, oxeye
Dandelion, common
Dandelion, false
Day flower
Deadnettle
Dichondra
Dock, broadleaf
Dock, curly
Doghane
Dogfennel
Dollarweed (pennywort)
Doveweed
Elderberry
Falseflax
False sunflower
Fiddleneck
Filaree, redstem
Filaree, whitestem
Fleabane, daisy (annual)
Fleabane, rough
Galinsoga, hairy
Galinsoga, smallflower
Goldenrod
Ground ivy
Groundsel
Gumweed
Hairy fleabane
Hawkweed, meadow
Hawkweed, mouseear
Hawkweed, orange
Hawkweed, yellow
Healall
Heartleaf drymary
Hemp
Henbit
Hoary cress
Horsenettle
Horseweed
Jimsonweed
Johnny-jumpup violet
Knawel
Knotweed, prostrate
Kochia
Lambsquarters
Lespedeza
Mallow, alkali
Mallow, bristly
Mallow, common
Mallow, dwarf
Mallow, Venice
Marcela
Matchweed
Mexicanweed
Moneywort
Morningglory, bigroot
Morningglory, ivyleaf
Morningglory, Mexican
Morningglory, red
Morningglory, tall
Morningglory, woolly
Mugwort
Mustard, ball
Mustard, black
Mustard, blue
Mustard, hedge
Mustard, Indian
Mustard, tansy
Mustard, tumble
Mustard, white
Mustard, wormseed
Nettle, stinging (burning)
Nettle, tall
Nettle, wood
Parsley-piert
Parsnip
Pearlwort
Pennycress (Frenchweed)
Pepperweed, perennial
Pepperweed, Virginia (peppergrass)
Pigweed, prostate
Pigweed, redroot
Pigweed, Russian
Pigweed, smooth
Pigweed, tumble
Pineappleweed
Plains coreopsis (tickseed)
Plantain, blackseed
Plantain, bracted
Plantain, broadleaf
Plantain, buckhorn
Plantain, hoary
Plantain, narrowleaf
Plantain, slender
Plantain, woolly
Poison ivy
Poison oak
Pokeweed
Prairie sunflower
Prickly lettuce (compass plant)
Prickly sida
Prostrate knotweed
Puncturevine
Purslane, common
Pusley, Brazil
Pusley, Florida
Pusley, large flower
Ragweed, bur
Ragweed, common
Ragweed, lanceleaf
Ragweed, western
Redstem filaree
Rough cinquefoil
Scarlet pimpernel
Shepherdspurse
Smartweed, ladysthumb
Smartweed, pale
Smartweed, Pennsylvania
Smooth chaff-flower
Smooth dock
Sorrel, red (sheep)
Sowthistle, annual
Sowthistle, spiny
Spanish needles
Speedwell, bi-lobed
Speedwell, common
Speedwell, corn
Speedwell, ivyleaf
Speedwell, Persian
Speedwell, purslane
Speedwell, slender
Speedwell, snow
Speedwell, thymeleaf
Spiny amaranth
Spurge, spotted
Spurge, prostrate
Spurweed
Star of Bethlehem
Strawberry, Indian mock
Tansy ragwort
Tanweed
Thistle, blessed
Thistle, bull
Thistle, Canada
Thistle, flodman (prairie)
Thistle, musk
Thistle, platte
Thistle, plumeless
Thistle, Russian
Thistle, Scotch
Thistle, wavyleaf
Thistle, yellow (horrible)
Trailing crownvetch
Velvetleaf
Vervain, blue
Vervain, hoary
Vervain, prostrate
Vervain, tall
Vetch, bird
Vetch, hairy
Vetch, milk
Vetch, narrowleaf
Vetch, wild
Virginia creeper
Western clematis
Western salsify
Wild aster
Wild buckweat
Wild carrot
Wild four-o-clock
Wild garlic
Wild lettuce
Wild marigold
Wild mustard (yellow rocket)
Wild onion
Wild parsnip
Wild radish
Wild rape
Wild strawberry
Wild sweet potato
Wild violet
Woodsorrel, creeping
Woodsorrel, violet
Woodsorrel, yellow (oxalis)
Woolly croton
Wormseed
Yarrow, common
Yarrow, western
Yarrow, yellow

GRASS WEEDS CONTROLLED/MALEZAS HERBOSAS CONTROLADAS: Crabgrass, large and smooth; Foxtail, green, yellow and giant; Barnyardgrass; Nutsedge, yellow

STOP READ THE ENTIRE LABEL BEFORE USE.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. *For outdoor residential use only.*

SHAKE WELL BEFORE USING.

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.

GENERAL INFORMATION:

Spectracide® Weed Stop® for Lawns plus Crabgrass Killer Concentrate is a selective herbicide that controls emerged broadleaf and grass weeds in established lawns. Symptoms you may expect to see on affected broadleaf weeds include leaf reddening, leaf and stem curling and twisting, chlorosis and necrosis. Symptoms expected on grass weeds include stunting, chlorosis, gradual reddening and necrosis.

WHERE TO USE

This product is intended for use by homeowners on residential lawns. This product should be applied only to the following species of turfgrass:

Cool season turfgrass	**Warm season turfgrass**
Kentucky bluegrass	Bermudagrass**
Perennial ryegrass	Zoysiagrass
Fescue spp., including tall, red and fine leaf fescues	Buffalograss

**Application to Bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to Bermudagrass when daytime temperatures exceed 85°F.

DO NOT USE this product on lawns containing bentgrass, St. Augustinegrass, bahiagrass, centipedegrass, carpetgrass. Also, do not use on dichondra, or lawns with desirable clovers or legumes. Do not use on vegetable gardens, fruits or ornamentals (flowers, trees, groundcovers, hedges, landscape/flower beds and shrubs).

WHEN TO USE

Apply this product to actively growing weeds during the early growth stages for the best results. Apply in spring or fall when daytime temperatures are between 45°F and 90°F. Fall applications may control winter annual and biennial weeds that may otherwise resprout the following spring. For heavy grass weed pressure, apply a second application at the same rate, 14 days after the first application.

CRABGRASS (large and smooth):

- **Spring:** For the best results, apply this product from crabgrass emergence until the six-leaf stage (two tillers) or before the crabgrass is 3 to 4 inches in height. Generally, this period begins or corresponds to the first time that you can see crabgrass in your lawn. Use two applications at the same rate with a 14-day interval for dense populations.

HOW MUCH TO USE

See the following table for calculating the correct amount of product to use for the area to be treated.
The maximum application rate is 6½ fl oz of product per 1,000 sq ft per application (0.6 lb 2,4-D acid equivalent per acre per application).
The maximum number of broadcast applications is limited to two per year.
The maximum seasonal rate is 13 fl oz of product per 1,000 sq ft (1.2 lb 2,4-D acid equivalent per acre), excluding spot treatments.

MEASUREMENT CALCULATIONS: Total lawn area in square feet = length x width.

EQUAL MEASURES:

1 fl oz = 2 Tbsp = 6 tsp	4 fl oz = ½ cup	16 fl oz = 1 pt
2 fl oz = 4 Tbsp	8 fl oz = 1 cup	32 fl oz = 1 qt

I. SPOT TREATMENT

To treat small areas or individual weeds use a pressure sprayer for better control. Mix the indicated amount of product listed with 1 gal of water for each 500 sq ft to be treated. Shake well before spraying. Spray the target weeds thoroughly. Nozzle adjustments may be necessary to deliver small spray droplets on the target weeds for uniform coverage.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION. Causes moderate eye irritation. Avoid contact with eyes or clothing. Harmful if swallowed. When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves.

Wash thoroughly with soap and water after handling. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

First Aid	
If in eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first five minutes, then continue rinsing eye. • Call a Poison Control Center or doctor for treatment advice.
If swallowed:	• Call a Poison Control Center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a Poison Control Center or doctor. • Do not give anything to an unconscious person.
Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment. You may also contact 1-800-917-5438 for emergency medical treatment advice.	

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.
Questions & Comments? Call 1-800-917-5438 or visit our website at www.spectracide.com

PRECAUCIONES

Riesgos para los seres humanos y los animales domésticos

PRECAUCIÓN. Causa irritación moderada a los ojos. Evite el contacto con los ojos o la ropa. Nocivo si se ingiere. Cuando utilice este producto, use camisa de manga larga, pantalones largos, calcetines, zapatos y guantes de goma/hule.

Después de manipular el producto, lávese muy bien con agua y jabón. Después de utilizar este producto, enjuague los guantes antes de quitárselos, quítese la ropa y lávela por separado antes de volver a usarla y rápidamente lávese bien las manos y la piel que estuvo expuesta, con agua y jabón. Quítese la ropa mojada y báñese en regadera tan pronto como sea posible.

Primeros Auxilios	
Si entra en contacto con los ojos:	• Mantenga los ojos abiertos y enjuague con agua lentamente, con cuidado, durante 15-20 minutos. Si hay lentes de contacto, retírelos después de los primeros 5 minutos, luego continúe enjuagando los ojos. • Llame a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento.
Si se ingiere:	• Llame inmediatamente a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento. • Si la persona puede tragar, haga que beba poco a poco un vaso de agua. • No induzca el vómito a menos que así lo indique un centro de control de envenenamientos o el médico. • No administre nada a una persona que haya perdido el conocimiento.
Cuando llame a un centro de control de envenenamientos o a un médico, o intente obtener tratamiento, tenga a la mano el envase o la etiqueta del producto. También puede llamar al 1-800-917-5438 para obtener información acerca del tratamiento médico de urgencia.	

NOTA: El comprador asume toda la responsabilidad en caso de no seguir las instrucciones sobre seguridad y modo de empleo.
¿Preguntas o comentarios? Llame al 1-800-917-5438.

Distributed by Spectrum Group, Div. of United Industries Corp.
PO Box 142642, St. Louis, MO 63114-0642
EPA Reg. No. 9688-268-8845 EPA Est. Nos. 39754-WI-1(R), 9688-MO-1(U), 79533-WI-1(BT), 39609-MO-1(BM) Circled letter is first letter of lot number. 17-12654 © 2008 UIC